



ROFLOXIN

INJECTABLE SOLUTION

COMPOSITION

Enrofloxacin.....	10,00 g
Excipient.....	100 ml

INDICATIONS

Treatment of bacterial infections caused by strains sensitive to enrofloxacin in cattle :

- Respiratory infections caused by *P. multocida*, *P. hemolytic* and *Mycoplasma bovis*.
- Digestive tract infections caused by *E. coli* and salmonella.
- Joint infections with *Mycoplasma bovis*.
- Treatment of acute mastitis with *E. coli*.

ADMINISTRATION AND DOSAGE

ROFLOXIN is intended to be administered by:

Subcutaneous use for the treatment of respiratory, digestive and joint infections at a dose of 5 mg of enrofloxacin per kg of bodyweight per day, or 5 ml of solution per 100 kg of bodyweight, for 3 to 5 days.

The intravenous route for the treatment of mastitis at a dose of 5 mg of enrofloxacin per kg of bodyweight per day, i.e. 5 ml of solution per 100 kg of bodyweight, for 2 consecutive days.

WITHDRAWAL PERIOD

Meat and offal:

Subcutaneous use: 21 days.

Intravenous use: 7 days.

Milk: 3 days or 6 milking .

STORAGE CONDITIONS

- 2 years in the original packaging, protected from light and heat.
- Stability after opening the container: 4 Weeks

PRESENTATIONS

50 ml and 100 ml bottles.