



# COMPOSITION

Amoxicillin (sf trihyrate)	10.00 g
Excipient qs	.100 g

# PHARMACEUTICAL PROPERTIES

Amoxicillin, active ingredient of ATLAMOX, is an antibiotic from the group of aminopenicillins widely used in veterinary medicine. Amoxicillin has bactericidal effects against many pathogenic germs in the target species. Amoxicillin has excellent pharmacokinetic properties: resistance in digestive acid pH, good digestive absorption and good tissue diffusion. The plasma half-life of amoxicillin is relatively short, resulting in a relatively short withdrawal period.

### TARGET SPECIES

Poultry and pre-ruminant calf.

# **INDICATIONS**

Infections with germs sensitive to amoxicillin in the target species, ATLAMOX is particularly indicated for the treatment of colibacillosis, salmonellosis, respiratory infections, infectious enteritis and staphylococcal disease.

### ADMINISTRATION AND DOSAGE

Oral use in drinking water or in the food ration.

Poultry: 1.00 g of powder / 10 kg bodyweight or a sachet of 100 g of ATLAMOX per 1000 kg once a day for 3 to 5 days.

The dose can be divided into two doses spaced 12 hours apart.

As an indication, ATLAMOX is to be used in drinking water at the rate of 1 sachet of 100 g / 100 liters of water or 1 sachet of 1 kg / ton of water.

Pre-ruminant calf: 1.00 g of powder / 10 kg / day for 3 to 5 days. The daily dose can be divided into two doses 12 hours apart. The product is to be administered after dilution in water or in milk replacer.



#### CONTRAINDICATIONS AND INTERACTIONS

- Hypersensitivity to Penicillins.
- Do not use in laying hens whose products are intended for human consumption.
- Do not use in rabbits and rodents.
- Do not combine with bacteriostatic antibacterial

#### SIDE EFFECTS

If the doses are respected, no side effects are observed in the target species.

#### PRECAUTIONS FOR USE

This soluble powder for oral preparation is intended to be dissolved in liquid food or water and cannot be used as it is. Respect the dose in dehydrated animals. Avoid handling this product in case of known allergy to amoxicillin. Handle this product with care to avoid any risk of exposure. Avoid breathing dust and avoid contact with skin and eyes.

#### WITHDRAWAL PERIOD

Meat and offal: 48 hours Eggs: Do not use in layers whose eggs or their derivatives are intended for human consumption.

#### **STORAGE CONDITIONS**

Keep the product in the original packaging, protect from light and at a temperature  $\leq 25$  ° C.

#### PHARMACEUTICAL FORM AND PRESENTATION

1 Kg sachets