



## COMPOSITION

Albendazole	. 2.50 g
Excipient qs	. 100 ml

## PHARMACEUTICAL PROPERTIES

Albendazole is an anthelmintic of the benzimidazole family endowed with a great activity on most internal parasites of ruminants, thus occupying an intermediate place between the anthelmintic endectocides, active on internal and external parasites, and other benzimidazoles no douvicides. ABZ is also active on adult forms of Fasciola hepatica. At a dose of 5 mg / kg, ABZ has an efficacy close to 80% on F. hepatica; at a double dose, it is greater than 90%. Note, however, that at this dose the activity is only 40% on immature stages.

### TARGET SPECIES

Cattle, sheep and goats.

# INDICATIONS

Treatment of gastrointestinal and pulmonary Strongyloses, Cestodoses and Fascioliasis with F. hepatica in ruminants

#### ADMINISTRATION AND DOSAGE

Oral use.

**Sheep and goats**: Gastrointestinal strongyloses, Dictyocaulosis and Tapeworm: 1.5 ml / 10 kg; Fascioliasis and other pulmonary strongyloses: 2 ml / 10 kg. **Cattle**: Gastrointestinal strongyloses, Dictyocaulosis and Tapeworm: 3 ml / 10 kg

### CONTRAINDICATIONS AND INTERACTIONS

Due to teratogenic and embryotoxic effects, dosages should be observed during the first month of gestation. Do not use in dairy females whose milk is intended for human consumption.



# SIDE EFFECTS

If the general conditions of use are respected, no undesirable effects are observed after treatment with Albendazole

### PRECAUTIONS FOR USE

- Respect the dose in females at the beginning of gestation.
- Shake well before use.
- Keep the product out of the reach of children

### WITHDRAWAL PERIOD

Meat and offal:

- Cattle: 14 days.
- Sheep and goats: 10 days

#### **STORAGE CONDITIONS**

Keep the product in the original packaging, protect from light and at a temperature  $\leq$  25  $^\circ$  C.

### PRESENTATIONS

Bottles of 100 ml, 250 ml, 500 ml, 1 liter and 2.5 L and 5-liter cans.