



COMPOSITION

Fosfomycin (s.f. calcium	n)> 20,00 g
Tylosin (s.f. tartrate)	5,00 g
Excipient q.s.p	100 g



PHARMACEUTICAL PROPERTIES

Fosfomycin is a bactericidal antibiotic produced by streptomyces. It has a broad spectrum encompassing both gram negative and gram-positive germs. It works by inhibiting the synthesis of the peptidoglycan precursors constituting the bacterial wall. In pharmacokinetics, fosfomycin is largely eliminated in the active form via the kidneys. Tylosin is one of the bacteriostatic macrolide antibiotics active mainly on gram positive bacteria and Mycoplasmas and on part of gram-negative bacteria (Pasteurella). The tylosin-fosfomycin combination is, in addition to its synergistic effect, endowed with a very broad spectrum.

TARGET SPECIES

Poultry.

INDICATIONS

TYLFOS is recommended for the treatment of gram-positive, gram-negative and mycoplasma infections in poultry.

ADMINISTRATION AND DOSAGE

Oral use in drinking water for 3 to 5 consecutive days at a dose of 160 mg powder / kg bodyweight.

CONTRAINDICATIONS AND INTERACTIONS

- Do not use in laying hens whose eggs or their derivatives are intended for human consumption.
- Positive interaction with beta lactam antibiotics, aminoglycosides, anti-staphylococcal agents.



SIDE EFFECTS

If the doses are respected, no side effects are observed in the target species.

PRECAUTIONS FOR USE

- This soluble powder for oral preparation is intended to be dissolved in liquid food or drinking water and cannot be used as it is.
- Respect the dose in dehydrated animals.
- Avoid handling this product in case of allergy Handle this product taking precautions to avoid any risk of exposure.
- Avoid breathing dust and avoid contact with skin and eyes.
- Keep the product out of the reach of children.

WITHDRAWAL PERIOD

- Meat and offal: 7 days
- Eggs: Do not use in laying hens whose eggs or their derivatives are intended for human consumption.

STORAGE CONDITIONS

- 2 years in the original packaging, protected from light and at a temperature below 25 ° C.
- Stability after opening the container: 3 Weeks Stability after dissolving: 6 to 8 hours

PRESENTATIONS:

1kg sachets