



# BOOSTER

DRINKABLE NUTRITIONAL SUPPLEMENT BASED ON VITAMINS AND AMINO ACIDS

## COMPOSITION

### VITAMINES

Vitamine A .....	10 MUI
Vitamine D3 .....	2 MUI
Nicotinamide (Vitamine B3) .....	16.25 g
D-Panthenol.....	7.50 g
Vitamine B12 .....	0.00125 g
Vitamine B2 .....	2.50 g
Acide folique.....	0.0716 g
Vitamine K3 .....	0.50 g
Vitamine B6 .....	1.125 g
Inositol .....	2.50 g
Biotine (Vitamine H) .....	0.001 g

### ACIDES AMINES

Alanine .....	11.5 g
Arginine .....	6.1 g
Acide aspartique .....	9.5 g
Phénylalanine .....	5.5 g
Cystéine.....	2.1 g
Acide glutamique .....	21.5 g
Glycine .....	9.1 g
Histidine .....	4.7 g
Isoleucine.....	6.7 g
Leucine .....	12.5 g
Lysine .....	9.5 g
DL-Méthionine .....	2.2 g
Proline .....	9.5 g
Sérine.....	7 g
Thréonine.....	5 g
Tyrosine.....	5.3g
Valine .....	5.2 g
Tryptophane .....	2 g
Excipient q.s.p.....	1000 ml

## TARGET SPECIES

Cattle, sheep, goats, camels, horses and poultry.

## INDICATIONS

• In deficiencies and nutritional disorders. Generally in weakness and recovering conditions.

In intensive productive cycles.

• During stress (heat stress, post-vaccination reaction, changes in food). • During the first days of life to increase the development of the digestive system. • During the moult of rabbits and poultry. • To increase lay, fertility and incubability in poultry and viability in chicks at hatching

When livestock are undernourished and convalescing. • During pregnancy. • During lactation. • To improve the yield and conversion index of cattle.

## ADMINISTRATION AND DOSAGE

Oral solution. Use in drinking water.

**Poultry** : 0.5ml to 1ml / liter of drinking water for 3 to 5 days.

- **In broilers** • On receipt of the

chicks: 1ml / liter for 3 to 5 days. • Week 3: 0.5ml/ liter for 2 to 3 days. •

Week 5: 0.5ml/ liter for 2 to 3 days.

- **In laying hens and breeders** : • On receipt of the chicks:

1ml/litre for 3 to 5 days. • Week 7-8: 0.5ml/ liter for 2 days. • At the start

of laying: Week 18 in layers: 1 ml / liter for 2 to 5 days. • Week 22 in

layers: 1 ml / liter for 2 to 5 days. • During the laying period: pulsatile doses of 0.5 ml/litre for 2 to 3 days per month.

**Other species:** 1 to 1.5 ml per 10 kg bodyweight, for 3 to 5 days in a row using a dosing gun or bottle.

**WITHDRAWAL PERIOD AND CONTRAINDICATIONS** : None.

## STORAGE CONDITIONS

- **Unopened container:** Keep the product in its commercial packaging, hermetically closed, stored in a cool, dry place away from light and at an ambient temperature < 25°C

- **After opening the bottle** :

## PHARMACEUTICAL FORM AND PRESENTATION

Drinkable solution.

Bottles of 100 ml, 250 ml, 500 ml, 1 liter, 2 liters and 5 liters.